

Trend-cycle Approach to Estimate Changes in Southern Canada's Water Yield

ROBERT BEMROSE

Statistics Canada

ENVIRONMENT ACCOUNTS AND STATISTICS DIVISION (EASD), CANADA

PETER MESZAROS y BENOIT QUENNEVILLE

Statistics Canada

BUSINESS SURVEY METHODS DIVISION (BSMD), CANADA

e-mail: Robby.Bemrose@statcan.gc.ca; Peter.Meszaros@statcan.gc.ca; Benoit.Quenneville@statcan.gc.ca

ABSTRACT

Quantifying how Canada's water yield has changed over time is an important component of the water accounts maintained by Statistics Canada. This study evaluates the movement in the series of annual water yield estimates for Southern Canada from 1971 to 2004. We estimated the movement in the series using a trend-cycle approach and found that water yield for Southern Canada has generally decreased over the period of observation.

Keywords: Trend-Cycle, Water Yield.

Un enfoque de ciclo-tendencia para estimar los cambios en el rendimiento del agua en Canadá

RESUMEN

La cuantificación de cómo ha evolucionado en el tiempo el rendimiento del agua en Canadá es un aspecto muy importante de la contabilidad del agua llevada a cabo por el Instituto de Estadística canadiense. Este artículo analiza los cambios en la serie de estimaciones del rendimiento anual de agua en el sur de Canadá desde 1971 hasta 2004. Así, estimamos los cambios en la serie mediante un enfoque de ciclo-tendencia, concluyendo que el rendimiento de agua en el sur de Canadá ha disminuido a lo largo del período estudiado.

Palabras clave: Ciclo-tendencia, rendimiento del agua.

Clasificación JEL: C30, C32.

Artículo recibido en agosto de 2010 y aceptado en septiembre de 2010.

Artículo disponible en versión electrónica en la página www.revista-eea.net, ref. 0-28311.

1. INTRODUCTION

Understanding how the amount of water yield¹ has changed in Canada is an integral part of Statistics Canada's Water Accounts. The framework for these accounts, the Canadian System of Environmental and Resource Accounts (CSERA), establishes an outline for estimates of stocks and flows of water moving within and between the natural environment and society. The framework offers an integrated view of water and its use by society through the compilation of environmental data that are consistent over space and time, national in coverage and compatible with economic accounts. An important part of the water accounting program is the measurement and analysis of these stocks and flows as they change over time. This is in keeping with one of Statistics Canada's objectives: the identification, measurement and tracking of trends in order to facilitate public debate and decision-making. This paper describes the trend estimation methodology used in the 2010 issue of Human Activity and the Environment: Freshwater supply and demand in Canada (Statistics Canada, 2010).

The search for trends in hydrometeorological data has become a regular undertaking given the ever-increasing need to understand how the magnitudes of present and historical components of the hydrological cycle evolve over time. Numerous studies have been conducted which use nonparametric rank-based techniques to determine the magnitude and significance of a trend found between time and streamflow. The Mann-Kendall test (Salmi *et al.*, 2002), one of the main methods used for trend estimation, provides a global robust estimate of the slope of the underlying trend in the series of annual water yield estimates. In this paper we present a methodology to complement this global estimate where the time series is deconstructed into several components separating the underlying trend-cycle from the irregularities in the series. To achieve this objective we first estimate a trend plus the cyclical component for the series of water yield estimates. We then describe the global trend over the span of the series by fitting a linear model to the trend-cycle estimate. To demonstrate this methodology, we used a water yield series from 1971 to 2004 for Southern Canada.


2. DATA AND METHODS

The water yield data consist of spatial estimates of average monthly unregulated flow (km^3) summarized annually across Canada from 1971 to 2004. These water

¹ The water yield is the amount of freshwater derived from unregulated streamflow (m^3/s) measurements for a given geographic area over a defined period of time. The freshwater streamflow is generated from a combination of baseflow, interflow, and overland flow originating from groundwater, precipitation and/or snowpack. The flow rate encompasses the hydrologic processes (for example, interception, infiltration, and evapotranspiration) and the state of water storage (for example, lakes, aquifers, snowpack, and soil moisture) within a drainage basin (Mosley and McKerchar, 1993); it is influenced by climatic and physiographic variables such as temperature and topography (Bemrose *et al.*, 2009).


yield estimates were produced by means of a geostatistical approach that involved kriging regionalized unregulated flow per unit area (km^3/km^2). After interpolation the km^3/km^2 estimates were converted back into km^3 for the desired accounting units. The unregulated flow data were obtained from the Water Survey of Canada's hydrometric database HYDAT (Environment Canada 2007). A detailed description of the methodology used to derive the water yield estimates can be found in Bemrose *et al.* (2009). Given the paucity of data, especially in the North, this study evaluated the changes in water yield for the area of 2,598,632 km^2 below the North-line² (Figure 1). The series of annual estimates of water yield below the North-line can be seen in Figure 2.

FIGURE 1
Statistics Canada's North-line in relation to the Canadian landmass.



² The North-line is based on a statistical area classification of the north by Statistics Canada reflecting a combination of sixteen social, biotic, economic and climatic characteristics that delineate north from south in Canada (McNiven and Puderer, 2000).

FIGURE 2
Average annual water yield for Southern Canada, 1971 to 2004.



A time series often consists of four components: the trend component which represents the systematic long-term movement over the span of the series; the cycle which describes the smooth movement around the trend; the seasonal component which consists of intra-year fluctuations, either monthly or quarterly; and the irregular (noise) component which is the residual not accounted for by the other components (Ladiray and Quenneville, 2001). Since the input series of water yield consisted of annual values, the seasonal component is not a component in the series. Consequently, the values were influenced only by the long-term trend, the quasi-cyclical fluctuations (the long-term trend and the quasi-cyclical fluctuations together are referred to as the signal), and the irregular component (*i.e.* the difference or ratio between the signal and the observations).

To estimate the signal, we used a nonparametric approach, whereby the series $\{X_t, t=1, 2, \dots, N\}$ was smoothed by a moving average of coefficients as follows:

$$\begin{aligned}\hat{X}_t &= \sum_{k=-p}^{+f} \theta_k X_{t+k} = \\ &= \theta_{-p} X_{t-p} + \dots + \theta_0 X_t + \dots + \theta_f X_{t+f}\end{aligned}\quad (1)$$

where $\theta_k, k = -p, \dots, f$ are the weights and θ_0 is the central weight for the moving averages from the p “past” and f “future” observations (Dagum, 1985; Ladiray and Quenneville, 2001). The resulting smoothed series is termed the trend-cycle because it consists of the long-term trend and cyclical fluctuations. The number of terms in the moving average is $p + f + 1$. When $p \neq f$ the filter is asymmetric; when $p = f$, and $\theta_{-k} = \theta_k, k = 1, \dots, p$ the filter is symmetric.

In this study, we used the Henderson (1916) filter to estimate the trend-cycle wherever there were enough observations to satisfy the p and f requirements. The Henderson filter reproduces local polynomials up to order 3. With a 17-term

Henderson filter we estimated the trend-cycle locally using 8 observations in the past and 8 observations in the future. To derive the 17 weights for the filter using Equation 2, we substituted 8 for n and the values were obtained for each k from -8 to 8 (Dagum *et al.*, 1996; Ladiray and Quenneville, 2001). The symmetric weights for the 17-term Henderson filter can be seen in Table 1.

$$\theta_k = \frac{315[(n-1)^2 - k^2][n^2 - k^2][(n+1)^2 - k^2][3n^2 - 16 - 11k^2]}{8n(n^2 - 1)(4n^2 - 1)(4n^2 - 9)(4n^2 - 25)} \quad (2)$$

Given the end-point problem with the Henderson symmetric filter, we used an asymmetric approach developed by Musgrave (1964) to determine the M coefficients $\{v_1, \dots, v_M\}$ using the symmetric N coefficients $\{w_1, \dots, w_N\}$ wherever $p \neq f$. Using the formula presented in Doherty (2001) and Findley *et al.* (1998) we calculated the Musgrave weights as follows:

$$v_j = w_j + \frac{1}{M} \sum_{i=M+1}^N w_i + \frac{\left(j - \frac{M+1}{2}\right)R}{1 + \frac{M(M-1)(M+1)}{12}R} \sum_{i=M+1}^N \left(i - \frac{M+1}{2}\right) w_i \quad (3)$$

where R was set to 4.5 for the 17-term Henderson filter. For a more detailed description of the Henderson filter and the associated Musgrave filter, see Ladiray and Quenneville (2001). The surrogate asymmetric coefficients used for the 17-term Henderson filter are presented in Table 1. The coefficients for each k from -8 to 8 sum to 1 for the Henderson filter when symmetric and asymmetric.

TABLE 1


Coefficients for the symmetric and asymmetric 17-term Henderson filter ($R = 4.5$).

The notation H_p_f indicates that the moving average has order $p+f+1$
with p points in the past and f points in the future.

k	H_{8_8}	H_{8_7}	H_{8_6}	H_{8_5}	H_{8_4}	H_{8_3}	H_{8_2}	H_{8_1}	H_{8_0}
-8	-0.00996	-0.00880	-0.00601	-0.00257	-0.00096	-0.00464	-0.01734	-0.04226	-0.08104
-7	-0.02037	-0.01945	-0.01727	-0.01466	-0.01347	-0.01608	-0.02468	-0.04041	-0.06242
-6	-0.01864	-0.01795	-0.01639	-0.01460	-0.01383	-0.01539	-0.01987	-0.02641	-0.03166
-5	0.00247	0.00291	0.00386	0.00483	0.00518	0.00469	0.00431	0.00697	0.01848
-4	0.04209	0.04230	0.04263	0.04278	0.04271	0.04328	0.04701	0.05886	0.08714
-3	0.09229	0.09227	0.09198	0.09131	0.09081	0.09244	0.10028	0.12133	0.16637
-2	0.14111	0.14085	0.13994	0.13845	0.13753	0.14023	0.15218	0.18241	0.24422
-1	0.17639	0.17589	0.17437	0.17205	0.17072	0.17447	0.19053	0.22996	0.30853
0	0.18923	0.18849	0.18636	0.18322	0.18146	0.18628	0.20644	0.25507	0.35040
1	0.17639	0.17541	0.17266	0.16871	0.16653	0.17240	0.19668	0.25449	0
2	0.14111	0.13989	0.13653	0.13175	0.12915	0.13609	0.16447	0	0
3	0.09229	0.09084	0.08686	0.08126	0.07824	0.08623	0	0	0
4	0.04209	0.04040	0.03580	0.02939	0.02594	0	0	0	0
5	0.00247	0.00054	-0.00468	-0.01191	0	0	0	0	0
6	-0.01864	-0.02081	-0.02664	0	0	0	0	0	0
7	-0.02037	-0.02278	0	0	0	0	0	0	0
8	-0.00996	0	0	0	0	0	0	0	0

The 17-term filter was selected for this series and for those presented in the Statistics Canada (2010) publication on the basis of its gain function displayed in Figure 3.

FIGURE 3
Gain function for 17-term symmetric Henderson filter.



The gain function shows the frequencies eliminated or preserved by the moving average (Ladiray and Quenneville, 2001). From Figure 3, it can be seen that the filter preserves 99% of the strength of cycles that repeat once every 34 years and 89% of those that repeat twice over 34 years; this is followed by a further decrease until all cycles that repeat more than 5 times are eliminated. For details on how the gain function is calculated, see Dagum *et al.* (1996) or Ladiray and Quenneville (2001).

The Henderson filter is available in seasonal adjustment software such as the Census X-11 (Shiskin *et al.*, 1967) and its variants including the X11-ARIMA (Dagum, 1988) and X12-ARIMA (Findley *et al.*, 1998). We implemented the latter X12-ARIMA in this study using the trend option. The software works through an iterative process where in the first stage it evaluates the trend-cycle for the original series (Figure 4) and determines the adjustment factors for the extreme values in the irregular component (Figure 5). In the second stage of processing the original series is modified for the extreme values, and the trend-cycle is recalculated (Figure 6); again, adjustment factors are determined for the extreme values in the irregular component (Figure 7). In the final stage of the processing chain, the original series is further modified for the extreme values and the final trend-cycle is recalculated (Figure 8). An overlay of the trend-cycles estimated through each stage of processing can be seen in Figure 9.

FIGURE 4
Original series of water yield and the first trend-cycle estimate.




FIGURE 5
Irregular component modified for extreme values of water yield.




FIGURE 6
Modified water yield and the second trend-cycle estimate compared to the original series of water yield.




FIGURE 7
Second adjustment of the irregular component modified for extreme values for water yield.






FIGURE 8

Second modified water yield and the final trend-cycle estimate compared to the original series of water yield.

**FIGURE 9**

Comparison of the first, second, and final trend-cycle estimate.



Local estimation of the trend-cycle and local down-weighting of extreme values is illustrated in Figures 7 and 8. Although the low water yield for 1978 is comparable to the levels observed in 1985, 1987, 1989, 1998, 2000, and 2001, it is considered atypical in comparison to its local neighbours. After producing the local trend-cycle for the series, we fitted a linear trend to the trend-cycle using the minimum least squares method to describe the global trend over the span of the series³.


³ Higher order polynomials could be fit. Adding higher orders to the global polynomial will increase the R^2 , which eventually will lead to a perfect fit to the trend-cycle.

3. RESULTS

Figure 10 displays the trend-cycle produced using the iterative process along with the linear fit to the trend-cycle, which describes the global trend. The intercept is estimated at 1388 km^3 , and the slope is $-3.52 \text{ km}^3/\text{yr}$.

FIGURE 10

Original water yield series, the final trend-cycle estimate, and the linear trend line.



The trend-cycle locally decreased from 1971 to 1987, increased until 1995, and then decreased again until 2004. Nineteen seventy-eight was the first year identified with an atypical water yield, but similar values have been observed in years since.

4. CONCLUSION

Water yield for the area below the North-line has globally decreased at a rate of $3.52 \text{ km}^3/\text{yr}$ from 1971 to 2004. Given that this is an aggregation of observations over a large and varied geographic area it is important to recognize that some regions may actually have had increasing water yields over the last 34 years, while the area as a whole experienced a decrease.

It is difficult to compare these results to those of other studies, given that most other studies focus on detecting changes that occur at localized areas over differing record lengths. However, some studies have made conclusions regarding the general trend for regions on the basis of an aggregation of clustered observations over fairly similar time periods. Zhang *et al.* (2001) found that annual average streamflow was for the most part decreasing across Southern Canada from 1967-1996. This finding is similar to that presented in Rivard *et al.* (2009), who

evaluated trends in groundwater recharge. They found that groundwater recharge has generally decreased for all observations below 55° latitude especially for Atlantic Canada from 1970 to 1997.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Estela Bee Dagum (Business Survey Methods Division, Time Series Research Analysis Centre) for providing valuable guidance through the steps in our analysis, and Dr. Soheil Rastan (Environment Accounts and Statistics Division) for helping us during the early stages of the project. We would also like to thank Mark Henry, Laura Kemp, and François Soulard (Environment Accounts and Statistics Division) for their work in the initial water yield study and their constructive review and edits of this manuscript.

REFERENCIAS BIBLIOGRÁFICAS

- BEMROSE, R.; KEMP, L.; HENRY, M. and SOULARD, F. (2009): The Water Yield for Canada as a Thirty-Year Average (1971 to 2000): Concepts, Methodology and Initial Results. *Environment Accounts and Statistics Analytical and Technical Paper Series*, Statistics Canada, Catalogue no. 16-001-MWE2009007.
- DAGUM, E.B.; CHHAB, N. and CHIU, K. (1996): Derivation and Analysis of the X11ARIMA and Census X11 Linear Filters. *Journal of Official Statistics*, vol. 12, no. 4, pp 329-347.
- DAGUM, E.B. (1985): *Moving Averages. Encyclopaedia of Statistical Sciences* (S. Kotz, and N. Johnson editors), John Wiley and Sons, vol. 5, pp. 630-634.
- DAGUM, E.B. (1988): *The X11ARIMA Seasonal Adjustment Method-Foundations and Users Manual*, Time Series Research and Analysis Centre, Statistics Canada.
- DOHERTY, M. (2001): The Surrogate Henderson Filters in X-11. *Australian and New Zealand Journal of Statistics*, vol. 43, no. 4, pp. 901-999.
- ENVIRONMENT CANADA (2007). National Data Archive, Hydrometric Database (HYDAT), http://www.wsc.ec.gc.ca/products/hydat/main_e.cfm?cname=hydat_iso_e.cfm (accessed January 7, 2009).
- FINDLEY, D.; MONSELL, B.; BELL, W.; OTTO, M. and CHEN, B. (1998): New Capabilities and Methods of the X12ARIMA Seasonal Adjustment Program. *Journal of Business Economic Statistics*, vol. 16, pp. 127-152.
- HENDERSON, R. (1916): Note on Graduation by Adjusted Average. *Transactions of the Actuarial Society of America*, vol. 17, pp. 43-48.
- McNIVEN, C. and PUDELER, H. (2000): Delineation of Canada's North: An Examination of the North-South Relationship in Canada. *Geography working Paper*, Statistics Canada, Catalogue no. 92F0138MIE.
- MOSLEY, P.M. and MCKERCHAR, A.I. (1993): Streamflow. In D.R. Maidment (Ed.), *Handbook of Hydrology* (pp. 8.1), McGraw-Hill, Inc.
- MUSGRAVE, J. (1964): A Set of End Weights to End all End Weights. Working Paper, US Bureau of Census, Washington.
- LADIRAY, D. and QUENNEVILLE, B. (2001): *Seasonal Adjustment with the X-11 Method*. Lecture Notes in Statistics, New York: Springer-Verlag, p. 225.

- RIVARD, C.; VIGNEAULT, H.; PIGGOTT, A.R.; LAROCQUE, M. and ANCTIL, F. (2009): Groundwater Recharge Trends in Canada. *Canadian Journal of Earth Sciences*, vol. 46, pp. 841-854.
- SALMI, T.; MÄÄTTÄ, A.; ANTILA, P.; RUOHO-AIROLA, T. and AMNELL, T. (2002): Detecting Trends of Annual Values of Atmospheric Pollutants by the Mann-Kendall Test and Sen's Slope Estimates – The Excel Template Application MAKESENS. Publication on Air Quality, *Finnish Meteorological Institute*, no. 31.
- SHISKIN, J.; YOUNG, A.H. and MUSGRAVE, J.C. (1967): The X-11 Variant of Census Method II Seasonal Adjustment, Technical Paper No. 15, U.S. *Bureau of the Census*.
- STATISTICS CANADA (2010): *Freshwater supply and demand in Canada, Human Activity and the Environment*, Catalogue no. 16-201-X.
- ZHANG, X.; HARVEY, K.D.; HOGG, W.D. and YUZYK, T.R. (2001): Trends in Canadian Streamflow. *Water Resource Research*, vol. 37, no. 4, pp. 987-998.